Author Index

Aaron, D.T., see Boswell, G.W. 513
Aasen, A.J., see Ruud-Christensen, M. 355
Aitio, A., see Pekari, K. 309
Alberti, K.G.M.M., see Cassidy, D.M. 1
Aldegunde, M., see Martin, F. 221
Allen, P.D., see Lema, M.J. 61
Ames, M.M.
— and Mathiesen, D.A.
High-performance liquid chromatographic assay and preclinical pharmacologic studies with the experimental antitumor agent batracylin 488
Arboix, M., see Guitart, R. 271
Baker, J.K., see Wall, G.M. 151
Baldew, G.S.
—, Volkers, K.J., De Goeij, J.J.M. and Vermeulen, N.P.E.
Determination of cisplatin and related platinum complexes in plasma ultrafiltrate and urine by high-performance liquid chromatography with on-line radioactivity detection 163
Barazi, S., see Marliac, Y. 461
Batt, A.M., see Schmitt, P. 495
Belolipetskaja, V.G.
—, Piotrovskii, V.K., Metelitsa, V.I. and Pavlinov, S.A.
Ion-exchange high-performance liquid chromatography in drug assay in biological fluids. V. Propranolol and metabolites 507
Bertolini, J.
—, Guthridge, M. and Hearn, M.T.W.
Rapid chromatographic isolation and immunoblot characterization of immunoreactive fibroblast growth factor-related polypeptides from various tissues 49
Bianchi, G.
High-performance liquid chromatographic assay of k-opioid selective benzeneacetamide derivatives (U50488, U69593 and PD117302) in rat plasma and brain 481
Bianchi, M., see Magnani, M. 215
Biondi, P.A.
—, Secchi, C., Negri, A., Tedeschi, G. and Ronchi, S.
Improved high-perfromance liquid chromatographic determination of diamine oxidase activity 209
Bochner, F., see Chen, Z.R. 367
Boer, E. de, see Teerlink, T. 418
Boswell, G.W.
—, Munoz, A.C., Aaron, D.T., Chin, C.L. and Quinn, K.P.
High-performance liquid chromatographic assay for RA 642, a compound with cardiovascular effect, in human and rabbit serum 513
Brandin, M.P., see Ruttyn, Y. 299
Brinkman, U.A.Th., see Irth, H. 321
Brune, K., see Geisslinger, G. 139
Brunnekreeft, J.W.I.
—, Eidhof, H. and Gerrits, J.
Optimized determination of thiochrome derivatives of thiamine and thiamine phosphates in whole blood by reversed-phase liquid chromatography with precolumn derivatization 89
Calvert, A.H., see Graham, M.A. 253
Canepa, A.
—, Perfumo, F., Carrea, A., Sanguineti, A., Piccardo, M.T. and Gusmano, R.
Measurement of free amino acids in polymorphonuclear leukocytes by high-performance liquid chromatograpy 200
Capizzi, R.L., see Wang, L.-M. 331
Carrea, A., see Canepa, A. 200
Cassidy, D.M.
—, Pratt, D.A., Taylor, R., Alberti, K.G.M.M. and Laker, M.F.
Capillary column gas chromatography-mass spectrometry for the determination of the fatty acid composition of human adipose tissue 1
Chen, Z.R.
—, Bochner, F. and Somogyi, A.
Simultaneous determination of codeine, norcodeine and morphine in biological fluids by high-performance liquid chromatography with fluorescence detection 367
Cheng, S.W.T.., see O'Connor, E.F. 240
Chin, C.L., see Boswell, G.W. 513
Chiou, S.-H.
— and Wang, K.-T.

Author Index

Aaron, D.T., see Boswell, G.W. 513
Aasen, A.J., see Ruud-Christensen, M. 355
Aitio, A., see Pekari, K. 309
Alberti, K.G.M.M., see Cassidy, D.M. 1
Aldegunde, M., see Martin, F. 221
Allen, P.D., see Lema, M.J. 61
Ames, M.M.
— and Mathiesen, D.A.
High-performance liquid chromatographic assay and preclinical pharmacologic studies with the experimental antitumor agent batracylin 488
Arboix, M., see Guitart, R. 271
Baker, J.K., see Wall, G.M. 151
Baldew, G.S.
—, Volkers, K.J., De Goeij, J.J.M. and Vermeulen, N.P.E.
Determination of cisplatin and related platinum complexes in plasma ultrafiltrate and urine by high-performance liquid chromatography with on-line radioactivity detection 163
Barazi, S., see Marliac, Y. 461
Batt, A.M., see Schmitt, P. 495
Belolipetskaja, V.G.
—, Piotrovskii, V.K., Metelitsa, V.I. and Pavlinov, S.A.
Ion-exchange high-performance liquid chromatography in drug assay in biological fluids. V. Propranolol and metabolites 507
Bertolini, J.
—, Guthridge, M. and Hearn, M.T.W.
Rapid chromatographic isolation and immunoblot characterization of immunoreactive fibroblast growth factor-related polypeptides from various tissues 49
Bianchi, G.
High-performance liquid chromatographic assay of k-opioid selective benzeneacetamide derivatives (U50488, U69593 and PD117302) in rat plasma and brain 481
Bianchi, M., see Magnani, M. 215
Biondi, P.A.
—, Secchi, C., Negri, A., Tedeschi, G. and Ronchi, S.
Improved high-perfromance liquid chromatographic determination of diamine oxidase activity 209
Bochner, F., see Chen, Z.R. 367
Boer, E. de, see Teerlink, T. 418
Boswell, G.W.
—, Munoz, A.C., Aaron, D.T., Chin, C.L. and Quinn, K.P.
High-performance liquid chromatographic assay for RA 642, a compound with cardiovascular effect, in human and rabbit serum 513
Brandin, M.P., see Ruttyn, Y. 299
Brinkman, U.A.Th., see Irth, H. 321
Brune, K., see Geisslinger, G. 139
Brunnekreeft, J.W.I.
—, Eidhof, H. and Gerrits, J.
Optimized determination of thiochrome derivatives of thiamine and thiamine phosphates in whole blood by reversed-phase liquid chromatography with precolumn derivatization 89
Calvert, A.H., see Graham, M.A. 253
Canepa, A.
—, Perfumo, F., Carrea, A., Sanguineti, A., Piccardo, M.T. and Gusmano, R.
Measurement of free amino acids in polymorphonuclear leukocytes by high-performance liquid chromatograpy 200
Capizzi, R.L., see Wang, L.-M. 331
Carrea, A., see Canepa, A. 200
Cassidy, D.M.
—, Pratt, D.A., Taylor, R., Alberti, K.G.M.M. and Laker, M.F.
Capillary column gas chromatography-mass spectrometry for the determination of the fatty acid composition of human adipose tissue 1
Chen, Z.R.
—, Bochner, F. and Somogyi, A.
Simultaneous determination of codeine, norcodeine and morphine in biological fluids by high-performance liquid chromatography with fluorescence detection 367
Cheng, S.W.T.., see O'Connor, E.F. 240
Chin, C.L., see Boswell, G.W. 513
Chiou, S.-H.
— and Wang, K.-T.

Peptide and protein hydrolysis by microwave irradiation 424
Choma, N., see Fukuda, E.K. 97
Cooper, J.K., see Srinivas, N.R. 262
Cucchiarini, L., see Magnani, M. 215
Dankers, J., see Macrae, P.V. 439
Davis, P.P., see Fukuda, E.K. 97
De Boer, E., see Teerlink, T. 418
Decouvelaere, B., see Schmitt, P. 495
De Goede, P.N.F.C., see De Kroon, I.F.I. 107
De Goeij, J.J.M., see Baldew, G.S. 163
De Graaf, S.S.N.
—, Riley-Stewart, C.A. and Evans, W.E. Improved high-performance liquid chromatographic method using loop-column extraction for analysis of idarubicin and idarubicinol in plasma 501
De Jong, G.J., see Irth, H. 321
De Kroon, I.F.I.
—, Langendijk, P.N.J. and De Goede, P.N.F.C. Simultaneous determination of midazolam and its three hydroxy metabolites in human plasma by electron-capture gas chromatography without derivatization 107
Dietzel, K., see Geisslinger, G. 139
D'Incalci, M., see Montaldo, P.G. 129
Discenza, C., see Selinger, K. 248
Duchateau, G.S.M.J.E., see Macrae, P.V. 439
Edwards, I.R., see Mason, R.W. 468
Eidhof, H., see Brunnekreeft, J.W.I. 89
Evans, W.E., see De Graaf, S.S.N. 501
Frei, R.W., see Irth, H. 321
Fukuda, E.K.
—, Choma, N. and Davis, P.P. Quantitation of the benzodiazepine antagonist flumazenil in human plasma by gas chromatography–mass spectrometry 97
Gaillot, J., see Schmitt, P. 495
Geisslinger, G.
—, Dietzel, K., Loew, D., Schuster, O., Rau, G., Lachmann, G. and Brune, K. High-performance liquid chromatographic determination of ibuprofen, its metabolites and enantiomers in biological fluids 139
Gerber, G., see Werner, A. 77
Gerrits, J., see Brunnekreeft, J.W.I. 89
Goede, P.N.F.C. de, see De Kroon, I.F.I. 107
Goeij, J.J.M. de, see Baldew, G.S. 163
Graaf, S.S.N. de, see De Graaf, S.S.N. 501
Graham, M.A.
—, Newell, D.R. and Calvert, A.H. Determination of the anthrapyrazole anticancer drug CI-941 in plasma and urine by solid-phase extraction and high-performance liquid chromatography 253
Guitart, R.
—, Puigdemont, A. and Arboix, M. Rapid headspace gas chromatographic method for the determination of liquid/gas partition coefficients 271
Gusmano, R., see Canepa, A. 200
Guthridge, M., see Bertolini, J. 49
Hanazono, H., see Iwata, T. 404
Hansen, S.H. Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
Haring, C., see Humpel, C. 235
Harvey, S.
—, Wiesler, D. and Novotny, M. Formation of cyclic enol ethers from a labile biological precursor: an example of analytical artifacts 27
Hashizume, Y., see Niwa, T. 397
Hayashi, T., see Tsuchiya, H. 291
Haymond, M.W., see Lecavalier, L. 410
Hearn, M.T.W., see Bertolini, J. 49
Hill, H.M., see Selinger, K. 248
Hirao, T., see Kikawa, Y. 193
Horber, F.F., see Lecavalier, L. 410
Hubbard, J.W., see Srinivas, N.R. 262
Humpel, C.
—, Haring, C. and Saria, A. Rapid and sensitive determination of clozapine in human plasma using high-performance liquid chromatography and amperometric detection 235
Iinuma, F., see Mawatari, K.-I. 389
Irth, H.
—, Tocklu, R., Welten, K., De Jong, G.J., Brinkman, U.A.Th. and Frei, R.W. Trace-level determination of 3′-azido-3′-deoxythymidine in human plasma by preconcentration on a silver(I)–thiol stationary phase with on-line reversed-phase

high-performance liquid chromatography 321
Iwata, T.
—, Hanazono, H., Yamaguchi, M., Nakamura, M. and Ohkura, Y.
Ultramicro determination of 7-dehydrocholesterol in rat skin by high-performance liquid chromatography with fluorescence detection 404
Jehl, F., see Koechlin, C. 379
Johansen, M., see Kurtzhals, P. 117
Jong, G.J. de, see Irth, H. 321
Kaneda, N., see Niwa, T. 397
Kikawa, Y.
—, Nakai, A., Shigematsu, Y., Sudo, M., Suehiro, Y. and Hirao, T.
Rapid and simle assay method to study the content and the biosynthesis of peptide leukotrienes in whole blood 193
Koechlin, C.
—, Jehl, F., Linger, L. and Monteil, H.
High-performance liquid chromatography for the determination of three new fluoroquinolones, fleroxacin, temafloxacin and A-64730, in biological fluids 379
Köppel, C.
—, Tenczer, J. and Wagemann, A.
Mass spectral characterization of urinary pipamperone metabolites and high-performance liquid chromatography assay for pipamperone plasma levels 432
Kowalewski, J., see Werner, A. 77
Krishna Murthy, D.V., see Ramesha, C.S. 37
Kroon, I.F.I. de, see De Kroon, I.F.I. 107
Kumps, A., see Wurth, C. 186
Kurtzhals, P.
—, Larsen, C. and Johansen, M.
High-performance size-exclusion chromatographic procedure for the determination of fluoresceinyl isothiocyanate dextrans of various molecular masses in biological media 117
Labadarios, D., see Shephard, G.S. 226
Lachmann, G., see Geisslinger, G. 139
Laker, M.F., see Cassidy, D.M. 1
Langendijk, P.N.J., see De Kroon, I.F.I. 107
Larsen, C., see Kurtzhals, P. 117
Lebel, G., see Selinger, K. 248
Lecavalier, L.
—, Horber, F.F. and Haymond, M.W.
Determination of plasma concentrations and ^{3}H-specific activity of phenylalanine in plasma using high-performance liquid chromatography 410
Lema, M.J.
—, Pluskal, M.G. and Allen, P.D.
High-performance ion-exchange chromatography of myosin using a DEAE-5PW column 61
Lho, D., see Shin, H. 448
Lindström, B., see Persson, K. 473
Linger, L., see Koechlin, C. 379
Loew, D., see Geisslinger, G. 139
Macrae, P.V.
—, Rance, D.J., Dankers, J. and Duchateau, G.S.M.J.E.
Analysis of a novel cardiac stimulant, UK-61 260, in human plasma by gas chromatography–mass spectrometry with selected-ion monitoring 439
Magnani, M.
—, Rossi, L., Bianchi, M., Cucchiarini, L. and Stocchi, V.
Reversed-phase liquid chromatographic determination of 2′,3′-dideoxycytidine in human blood samples 215
Mallet, A.I., see Murray, S. 15
Mardens, Y., see Wurth, C. 186
Marliac, Y.
— and Barazi, S.
Dosage du nitrazépam inchangé dans le plasma par chromatographie en phase gazeuse 461
Martin, F.
— and Aldegunde, M.
Simple high-performance liquid chromatographic method with electrochemical detection for the determination of indoleamines in tissue and plasma 221
Mason, R.W.
— and Edwards, I.R.
High-performance liquid chromatographic determination of rhodamine B in rabbit and human plasma 468
Mathiesen, D.A., see Ames, M.M. 488
Matsuo, N., see Takeyama, N. 69
Mawatari, K.-I.
—, Iinuma, F. and Watanabe, M.
Fluorimetric determination of indomethacin in human serum by high-perform-

ance liquid chromatography coupled with post-column photochemical reaction with hydrogen peroxide 389
Metelitsa, V.I., see Belolipetskaja, V.G. 507
Midha, K.K., see Srinivas, N.R. 262
Montaldo, P.G.
— and D'Incalci, M.
High-performance liquid chromatographic determination of 5-bromodeoxyuridine in human plasma 129
Monteil, H., see Koechlin, C. 379
Moore, C.M.
— and Oliver, J.S.
Rapid extraction and determination of xylazine in greyhound urine using high-performance liquid chromatography 519
Munoz, A.C., see Boswell, G.W. 513
Murray, S.
—, O'Mally, G., Taylor, I.K., Mallet, A.I. and Taylor, G.W.
Assay for N^{τ}-methylimidazoleacetic acid, a major metabolite of histamine, in urine and plasma using capillary column gas chromatography–negative-ion mass spectrometry 15
Nagatsu, T., see Niwa, T. 397
Nakai, A., see Kikawa, Y. 193
Nakamura, M., see Iwata, T. 404
Negri, A., see Biondi, P.A. 209
Newell, D.R., see Graham, M.A. 253
Niwa, T.
—, Takeda, N., Sasaoka, T., Kaneda, N., Hashizume, Y., Yoshizumi, H., Tatematsu, A. and Nagatsu, T.
Detection of tetrahydroisoquinoline in parkinsonian brain as an endogenous amine by use of gas chromatography–mass spectrometry 397
North, W.G., see O'Connor, E.F. 240
Novotny, M., see Harvey, S. 27
O'Connor, E.F.
—, Cheng, S.W.T. and North, W.G.
Simultaneous extraction and chromatographic analysis of morphine, dilaudid, naltrexone and naloxone in biological fluids by high-performance liquid chromatography with electrochemical detection 240
Ohkura, Y., see Iwata, T. 404
Oliver, J.S., see Moore, C.M. 519
O'Mally, G., see Murray, S. 15
Park, J., see Shin, H. 448
Pavlinov, S.A., see Belolipetskaja, V.G. 507
Pekari, K.
—, Riekkola, M.-L. and Aitio, A.
Simultaneous determination of benzene and toluene in the blood using head-space gas chromatography 309
Perfumo, F., see Canepa, A. 200
Persson, K.
—, Lindström, B., Spalding, D., Wahlström, A. and Rane, A.
Determination of codeine and its metabolites in human blood plasma and in microsomal incubates by high-performance liquid chromatography with ultraviolet detection 473
Piccardo, M.T., see Canepa, A. 200
Pickett, W.C., see Ramesha, C.S. 37
Piotrovskii, V.K., see Belolipetskaja, V.G. 507
Pluskal, M.G., see Lema, M.J. 61
Porsius, A.J., see Verrijk, R. 265
Pratt, D.A., see Cassidy, D.M. 1
Puigdemont, A., see Guitart, R. 271
Quinn, K.P., see Boswell, G.W. 513
Qureshi, A.R., see Qureshi, G.A. 281
Qureshi, G.A.
— and Qureshi, A.R.
Determination of free amino acids in biological samples: problems of quantitation 281
Rabenstein, D.L., see Yamashita, G.T. 341
Ramesha, C.S.
—, Pickett, W.C. and Krishna Murthy, D.V.
Sensitive method for the analysis of phospholipid subclasses and molecular species as 1-anthroyl derivatives of their diglycerides 37
Rance, D.J., see Macrae, P.V. 439
Rane, A., see Persson, K. 473
Rasmussen, K.E., see Ruud-Christensen, M. 355
Rau, G., see Geisslinger, G. 139
Riekkola, M.-L., see Pekari, K. 309
Riley-Stewart, C.A., see De Graaf, S.S.N. 501
Ronchi, S., see Biondi, P.A. 209
Rooij, H.H. van, see Verrijk, R. 265
Rossi, L., see Magnani, M. 215
Ruttyn, Y.
—, Brandin, M.P. and Vijayalakshmi, M.A.
Chromatography of human plasma on

aminohexyl Sepharose: separation of factor VIII/vWf and behaviour of factors II, VII, IX and X and antithrombin III 299
Ruud-Christensen, M.
—, Aasen, A.J., Rasmussen, K.E. and Salvesen, B.
High-performance liquid chromatographic determination of (*R*)- and (*S*)-proxyphylline in human plasma 355
Salvesen, B., see Ruud-Christensen, M. 355
Sanguineti, A., see Canepa, A. 200
Saria, A., see Humpel, C. 235
Sasaoka, T., see Niwa, T. 397
Schmitt, P.
—, Batt, A.M., Decouvelaere, B., Gaillot, J. and Siest, G.
Preparative high-performance liquid chromatographic method for the 2-hydroxy metabolite of carpipramine, excreted by isolated perfused liver 495
Schuster, O., see Geisslinger, G. 139
Secchi, C., see Biondi, P.A. 209
Selinger, K.
—, Lebel, G., Hill, H.M. and Discenza, C.
High-performance liquid chromatographic method for the analysis of benztropine in human plasma 248
Shephard, G.S.
—, Van der Westhuizen, L. and Labadarios, D.
Analysis of vitamin B_6 vitamers in human tissue by cation-exchange high-performance liquid chromatography 226
Shigematsu, Y., see Kikawa, Y. 193
Shin, H.
—, Lho, D. and Park, J.
Simultaneous determination of propoxyphene and norpropoxyphene in biological samples by gas chromatography using an on-column injection technique with a fused-silica capillary column 448
Siems, W., see Werner, A. 77
Siest, G., see Schmitt, P. 495
Somogyi, A, see Chen, Z.R. 367
Spalding, D., see Persson, K. 473
Srinivas, N.R.
—, Cooper, J.K., Hubbard, J.W. and Midha, K.K.
Isothermal capillary gas chromatography with electron-capture detection of heptafluorobutyryl-L-prolyl derivatives of chiral amphetamines 262
Stocchi, V., see Magnani, M. 215
Sudo, M., see Kikawa, Y. 193
Suehiro, Y., see Kikawa, Y. 193
Sunzel, M.
Determination of midazolam and the α-hydroxy metabolite by gas chromatography in small plasma volumes 455
Takagi, D., see Takeyama, N. 69
Takeda, N., see Niwa, T. 397
Takeyama, N.
—, Matsuo, N., Takagi, D. and Tanaka, T.
Determination of overt carnitine palmitoyltransferase by reversed-phase high-performance liquid chromatography 69
Tanaka, T, see Takeyama, N. 69
Tatematsu, A., see Niwa, T. 397
Taylor, G.W., see Murray, S. 15
Taylor, I.K., see Murray, S. 15
Taylor, R., see Cassidy, D.M. 1
Tedeschi, G., see Biondi, P.A. 209
Teerlink, T.
— and De Boer, E.
Determination of 3-methylhistidine in urine by high-performance liquid chromatography using pre-column derivatization with 9-fluorenylmethyl chloroformate 418
Tenczer, J., see Köppel, C. 432
Tocklu, R., see Irth, H. 321
Tsuchiya, H.
—and Hayashi, T.
Determination of L-3,4-dihydroxyphenylalanine in blood by high-performance liquid chromatography after solvent extraction 291
Van der Westhuizen, L., see Shephard, G.S. 226
Van Rooij, H.H., see Verrijk, R. 265
Vermeulen, N.P.E., see Baldew, G.S. 163
Verrijk, R.
—, Van Rooij, H.H., Wemer, H. and Porsius, A.J.
High-performance liquid chromatographic determination of milrinone in biological tissues and fluids 265
Vijayalakshmi, M.A., see Ruttyn, Y. 299
Volkers, K.J., see Baldew, G.S. 163
Wagemann, A., see Köppel, C. 432
Wahlström, A., see Persson, K. 473
Wall, G.M.
— and Baker, J.K.
Determination of baclofen and α-baclofen

in rat liver homogenate and human urine using solid-phase extraction, *o*-phthalaldehyde–*tert.*-butyl thiol derivatization and high-performance liquid chromatography with amperometric detection 151
Wang, K.-T., see Chiou, S.-H. 424
Wang, L.-M.
—, Woodward, C.N., White, J.C. and Capizzi, R.L.
Determination of ^{3}H-labelled cytosine arabinoside and eight metabolites in cell extracts by high-performance liquid chromatography and scintillation spectrometry 331
Watanabe, M., see Mawatari, K.-I. 389
Welten, K., see Irth, H. 321
Wemer, J., see Verrijk, R. 265
Werner, A.
—, Siems, W., Kowalewski, J. and Gerber, G.
Interrelationships between purine nucleotide degradation and radical formation during intestinal ischaemia and reperfusion: an application of ion-pair high-performance liquid chromatography of nucleotides, nucleosides and nucleobases 77
Westhuizen, L. van der, see Shephard, G.S. 226
White, J.C., see Wang, L.-M. 331
Wiesler, D., see Harvey, S. 27
Woodward, C.N., see Wang, L.-M. 331
Wurth, C.
—, Kumps, A. and Mardens, Y.
Urinary organic acids: retention indices on two capillary gas chromatography columns 186
Yamaguchi, M., see Iwata, T. 404
Yamashita, G.T.
— and Rabenstein, D.L.
Determination of penicillamine, penicillamine disulfide and penicillamine–glutathione mixed disulfide by high-performance liquid chromatography with electrochemical detection 341
Yoshizumi, H., see Niwa, T. 397

Subject Index

A-64730
HPLC for the determination of three new fluoroquinolones, fleroxacin, temafloxacin and A-64730 in biological fluids 379
N-Acetyl-5-aminosalicylic acid
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
N-Acetylhydroxysulphapyridine
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
N-Acetylsulphapyridine
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
N-Acetylsulphapyridine O-glucuronide
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
Amines
Detection of tetrahydroisoquinoline in parkinsonian brain as an endogenous amine by use of GC–MS 397
Amino acids
Measurement of free amino acids in polymorphonuclear leukocytes by HPLC 200
Amino acids
Determination of free amino acids in biological samples: problems of quantitation 281
Amino acids
Peptide and protein hydrolysis by microwave irradiation 424
5-Aminosalicylic acid
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
Amphetamines
Isothermal capillary GC with electron-capture detection of heptafluorobutyryl-L-prolyl derivatives of chiral amphetamines 262
Anthrapyrazole
Determination of the anthrapyrazole anticancer drug CI-941 in plasma and urine by solid-phase extraction and HPLC 253
Antithrombin III
Chromatography of human plasma on aminohexyl Sepharose: separation of factor VIII/vWf and behaviour of factors II, VII, IX and X and antithrombin III 299
3′-Azido-3′-deoxythymidine
Trace-level determination of 3′-azido-3′-deoxythymidine in human plasma by preconcentration on a silver(I)–thiol stationary phase with on-line reversed-phase HPLC 321
Baclofen
Determination of baclofen and α-baclofen in rat liver homogenate and human urine using solid-phase extraction, *o*-phthalaldehyde–*tert.*-butyl thiol derivatization and HPLC with amperometric detection 151
Batracylin
HPLC assay and preclinical pharmacologic studies with the experimental antitumor agent batracylin 488
Benzene
Simultaneous determination of benzene and toluene in the blood using head-space GC 309
Benzeneacetamide derivatives, k-opioid selective
HPLC assay of k-opioid selective benzeneacetamide derivatives (U50488, U69593 and PD117302) in rat plasma and brain 481
Benzodiazepine antagonist
Quantitation of the benzodiazepine antagonist flumazenil in human plasma by GC–MS 97
Benztropine
HPLC method for the analysis of benztropine in human plasma 248

2,2′-{[4,8-Bis(diethylamino)pyrimido[5,4-*d*]pyrimidine-2,6-diyl]bis[2-methoxyethyl)imino]}bisethanol
HPLC assay for RA 642, a compound with cardiovascular effect, in human and rabbit serum 513
5-Bromodeoxyuridine
HPLC determination of 5-bromodeoxyuridine in human plasma 129
Carboxyibuprofen
HPLC determination of ibuprofen, its metabolites and enantiomers in biological fluids 139
Carnitine palmitoyltransferase
Determination of overt carnitine palmitoyltransferase by reversed-phase HPLC 69
Carpipramine
Preparative HPLC method for the 2-hydroxy metabolite of carpipramine, excreted by isolated perfused liver 495
CI-941
Determination of the anthrapyrazole anticancer drug CI-941 in plasma and urine by solid-phase extraction and HPLC 253
Cisplatin
Determination of cisplatin and related platinum complexes in plasma ultrafiltrate and urine by HPLC with on-line radioactivity detection 163
Clotting factors
Chromatography of human plasma on aminohexyl Sepharose: separation of factor VIII/vWf and behaviour of factors II, VII, IX and X and antithrombin III 299
Clozapine
Rapid and sensitive determination of clozapine in human plasma using HPLC and amperometric detection 235
Codeine
Simultaneous determination of codeine, norcodeine and morphine in biological fluids by HPLC with fluorescence detection 367
Codeine
Determination of codeine and its metabolites in human blood plasma and in microsomal incubates by HPLC with UV detection 473
Codeine-6-glucuronide
Determination of codeine and its metabolites in human blood plasma and in microsomal incubates by HPLC with UV detection 473
Cyclic enol ethers
Formation of cyclic enol ethers from a labile biological precursor: an example of analytical artifacts 27
Cytosine arabinoside
Determination of ^{3}H-labelled cytosine arabinoside and eight metabolites in cell extracts by HPLC and scintillation spectrometry 331
Cytosine arabinoside phosphates
Determination of ^{3}H-labelled cytosine arabinoside and eight metabolites in cell extracts by HPLC and scintillation spectrometry 331
7-Dehydrocholesterol
Ultramicro determination of 7-dehydrocholesterol in rat skin by HPLC with fluorescence detection 404
Dextrans
High-performance size-exclusion chromatographic procedure for the determination of fluoresceinyl isothiocyanate dextrans of various molecular masses in biological media 117
Diamine oxidase
Improved HPLC determination of diamine oxidase activity 209
2′,3′-Dideoxycytidine
Reversed-phase LC determination of 2′,3′-dideoxycytidine in human blood samples 215
L-3,4-Dihydroxyphenylalanine
Determination of L-3,4-dihydroxyphenylalanine in blood by HPLC after solvent extraction 291
Dilaudid
Simultaneous extraction and chromatographic analysis of morphine, dilaudid, naltrexone and naloxone in biological fluids by HPLC with electrochemical detection 240
6-(2,4-Dimethyl-1H-imidazol-l-yl)-8-methyl-2(1H)-quinolinone
Analysis of a novel cardiac stimulant, UK-61 260, in human plasma by GC–MS with selected-ion monitoring 439

Enol ethers
Formation of cyclic enol ethers from a labile biological precursor: an example of analytical artifacts 27
Enzymes
Determinatiuon of overt carnitine palmitoyltransferase by reversed-phase HPLC 69
Enzymes
Improved HPLC determination of diamine oxidase activity 209
Factors II, VII, IX and X
Chromatography of human plasma on aminohexyl Sepharose: separation of factor VIII/vWf and behaviour of factors II, VII, IX and X and antithrombin III 299
Factor VIII/vWf
Chromatography of human plasma on aminohexyl Sepharose: separation of factor VIII/vWf and behaviour of factors II, VII, IX and X and antithrombin III 299
Fatty acids
Capillary column GC–MS for the determination of the fatty acid composition of human adipose tissue 1
Fibroblast growth factor
Rapid chromatographic isolation and immunoblot characterization of immunoreactive fibroblast growth factor-related polypeptides from various tissues 49
Fleroxacin
HPLC for the determination of three new fluoroquinolones, fleroxacin, temafloxacin and A-64730 in biological fluids 379
Flumazenil
Quantitation of the benzodiazepine antagonist flumazenil in human plasma by GC–MS 97
Fluoresceinyl isothiocyanate dextrans
High-performance size-exclusion chromatographic procedure for the determination of fluoresceinyl isothiocyanate dextrans of various molecular masses in biological media 117
Fluoroquinolones
HPLC for the determination of three new fluoroquinolones, fleroxacin, temafloxacin and A-64730 in biological fluids 379
Histamine
Assay for N^{τ}-methylimidazoleacetic acid, a major metabolite of histamine, in urine and plasma using capillary column GC–negative-ion MS 15
2-Hydroxycarpipramine
Preparative HPLC method for the 2-hydroxy metabolite of carpipramine, excreted by isolated perfused liver 495
Hydroxyibuprofen
HPLC determination of ibuprofen, its metabolites and enantiomers in biological fluids 139
5-Hydroxy-3-indoleacetic acid
Simple HPLC method with electrochemical detection for the determination of indoleamines in tissue and plasma 221
6-Hydroxy-6-methyl-3-heptanone
Formation of cyclic enol ethers from a labile biological precursor: an example of analytical artifacts 27
1-Hydroxymethyl-4-hydroxymidazolam
Simultaneous determination of midazolam and its three hydroxy metabolites in human plasma by electron-capture GC without derivatization 107
1-Hydroxymethylmidazolam
Simultaneous determination of midazolam and its three hydroxy metabolites in human plasma by electron-capture GC without derivatization 107
4-Hydroxymidazolam
Simultaneous determination of midazolam and its three hydroxy metabolites in human plasma by electron-capture GC without derivatization 107
α-Hydroxymidazolam
Determination of midazolam and the α-hydroxy metabolite by GC in small plasma volumes 455
4-Hydroxypropranolol
Ion-exchange HPLC in drug assay in biological fluids. V. Propranolol and metabolites 507
Hydroxysulphapyridine
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
5-Hydroxytryptamine
Simple HPLC method with electrochemical detection for the determination of indoleamines in tissue and plasma 221

5-Hydroxytryptophan
Simple HPLC method with electrochemical detection for the determination of indoleamines in tissue and plasma 221
Ibuprofen
HPLC determination of ibuprofen, its metabolites and enantiomers in biological fluids 139
Idarubicin
Improved HPLC method using loop–column extraction for analysis of idarubicin and idarubicinol in plasma 501
Idarubicinol
Improved HPLC method using loop–column extraction for analysis of idarubicin and idarubicinol in plasma 501
Indoleamines
Simple HPLC method with electrochemical detection for the determination of indoleamines in tissue and plasma 221
Indomethacin
Fluorimetric determination of indomethacin in human serum by HPLC coupled with post-column photochemical reaction with hydrogen peroxide 389
Leukotrienes
Rapid and simple assay method to study the content and the biosynthesis of peptide leukotrienes in whole blood 193
Liquid/gas partition coefficients
Rapid head space GC method for the determination of liquid/gas partition coefficients 271
3-Methylhistidine
Determination of 3-methylhistidine in urine by HPLC using pre-column derivatization with 9-fluorenylmethyl chloroformate 418
N^{τ}-Methylimidazoleacetic acid
Assay for N^{τ}-methylimidazoleacetic acid, a major metabolite of histamine, in urine and plasma using capillary column GC–negative-ion MS 15
Microwave irradiation
Peptide and protein hydrolysis by microwave irradiation 424
Midazolam
Simultaneous determination of midazolam and its three hydroxy metabolites in human plasma by electron-capture GC without derivatization 107
Midazolam
Determination of midazolam and the α-hydroxy metabolite by GC in small plasma volumes 455
Milrinone
HPLC determination of milrinone in biological tissues and fluids 265
Morphine
Simultaneous extraction and chromatographic analysis of morphine, dilaudid, naltrexone and naloxone in biological fluids by HPLC with electrochemical detection 240
Morphine
Simultaneous determination of codeine, norcodeine and morphine in biological fluids by HPLC with fluorescence detection 367
Morphine
Determination of codeine and its metabolites in human blood plasma and in microsomal incubates by HPLC with UV detection 473
Myosin
High-performance ion-exchange chromatography of myosin using a DEAE-5PW column 61
Naloxone
Simultaneous extraction and chromatographic analysis of morphine, dilaudid, naltrexone and naloxone in biological fluids by HPLC with electrochemical detection 240
Naltrexone
Simultaneous extraction and chromatographic analysis of morphine, dilaudid, naltrexone and naloxone in biological fluids by HPLC with electrochemical detection 240
Nitrazepam
Determination of intact nitrazepam in plasma by GC 461
Norcodeine
Simultaneous determination of codeine, norcodeine and morphine in biological fluids by HPLC with fluorescence detection 367
Norcodeine
Determination of codeine and its metabolites in human blood plasma and in mi-

crosomal incubates by HPLC with UV detection 473
Norpropoxyphene
Simultaneous determination of propoxyphene and norpropoxyphene in biological samples by GC using an on-column injection technique with a fused-silica capillary column 448
Nucleobases
Interrelationships between purine nucleotide degradation and radical formation during intestinal ischaemia and reperfusion: an application of ion-pair HPLC of nucleotides, nucleosides and nucleobases 77
Nucleosides
Interrelationships between purine nucleotide degradation and radical formation during intestinal ischaemia and reperfusion: an application of ion-pair HPLC of nucleotides, nucleosides and nucleobases 77
Nucleotides
Interrelationships between purine nucleotide degradation and radical formation during intestinal ischaemia and reperfusion: an application of ion-pair HPLC of nucleotides, nucleosides and nucleobases 77
Nucleotides
Reversed-phase LC determination of 2′,3′-dideoxycytidine in human blood samples 215
Organic acids
Urinary organic acids: retention indices on two capillary GC columns 186
Partition coefficients
Rapid head space GC method for the determination of liquid/gas partition coefficients 271
PD117302
HPLC assay of k-opioid selective benzeneacetamide derivatives (U50488, U69593 and PD117302) in rat plasma and brain 481
Penicillamine
Determination of penicllamine, penicillamine disulfide and penicillamine–glutathione mixed disulfide by HPLC with electrochemical detection 341
Penicillamine disulfide
Determination of penicillamine, penicillamine disulfide and penicillamine–glutathione mixed disulfide by HPLC with electrochemical detection 341
Penicillamine–glutathione mixed disulfide
Determination of penicillamine, penicillamine disulfide and penicillamine–glutathione mixed disulfide by HPLC with electrochemical detection 341
Peptide leukotrienes
Rapid and simple assay method to study the content and the biosynthesis of peptide leukotrienes in whole blood 193
Peptides
Peptide and protein hydrolysis by microwave irradiation 424
Phenylalanine
Determination of plasma concentrations and ^{3}H-specific activity of phenylalanine in plasma using HPLC 410
Phospholipids
Sensitive method for the analysis of phospholipid subclasses and molecular species as 1-anthroyl derivatives of their diglycerides 37
Pipamperone
Mass spectral characterization of urinary pipamperone metabolites and HPLC assay for pipamperone plasma levels 432
Platinum complexes
Determination of cisplatin and related platinum complexes in plasma ultrafiltrate and urine by HPLC with on-line radioactivity detection 163
Polypeptides
Rapid chromatographic isolation and immunoblot characterization of immunoreactive fibroblast growth factor-related polypeptides from various tissues 49
Propoxyphene
Simultaneous determination of propoxyphene and norpropoxyphene in biological samples by GC using an on-column injection technique with a fused-silica capillary column 448
Propranolol
Ion-exchange HPLC in drug assay in biological fluids. V. Propranolol and metabolites 507

Propranolol glucuronide
Ion-exchange HPLC in drug assay in biological fluids. V. Propranolol and metabolites 507
Proteins
High-performance ion-exchange chromatography of myosin using a DEAE-5PW column 61
Proteins
Peptide and protein hydrolysis by microwave irradiation 424
Proxyphylline
HPLC determination of (*R*)- and (*S*)-proxyphylline in human plasma 355
Purine nucleotides
Interrelationships between purine nucleotide degradation and radical formation during intestinal ischaemia and reperfusion: an application of ion-pair HPLC of nucleotides, nucleosides and nucleobases 77
Pyridoxal
Analysis of vitamin B_6 vitamers in human tissue by cation-exchange HPLC 226
Pyridoxamine
Analysis of vitamin B_6 vitamers in human tissue by cation-exchange HPLC 226
Pyridoxic acid
Analysis of vitamin B_6 vitamers in human tissue by cation-exchange HPLC 226
Pyridoxine
Analysis of vitamin B_6 vitamers in human tissue by cation-exchange HPLC 226
RA 642
HPLC assay for RA 642, a compound with cariovascular effect, in human and rabbit serum 513
Rhodamine B
HPLC determination of rhodamine B in rabbit and human plasma 468
Sulphapyridine
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
Sulphapyridine O-glucuronide
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
Sulphasalazine
Simple and rapid method for the simultaneous determination of the eight main metabolites and conjugates of sulphasalazine in human plasma, urine and faeces using dynamically modified silica 175
Steroids
Ultramicro determination of 7-dehydrocholesterol in rat skin by HPLC with fluorescence detection 404
Temafloxacin
HPLC for the determination of three new fluoroquinolones, fleroxacin, temafloxacin and A-64730 in biological fluids 379
Tetrahydroisoquinoline
Detection of tetrahydroisoquinoline in parkinsonian brain as an endogenous amine by use of GC–MS 397
Thiamine
Optimized determination of thiochrome derivatives of thiamine and thiamine phosphates in whole blood by reversed-phase LC with precolumn derivatization 89
Thiamine phosphates
Optimized determination of thiochrome derivatives of thiamine and thiamine phosphates in whole blood by reversed-phase LC with precolumn derivatization 89
Toluene
Simultaneous determination of benzene and toluene in the blood using head-space GC 309
U50488
HPLC assay of k-opioid selective benzeneacetamide derivatives (U50488, U69593 and PD117302) in rat plasma and brain 481
U69593
HPLC assay of k-opioid selective benzeneacetamide derivatives (U50488, U69593 and PD117302) in rat plasma and brain 481
UK-61 260
Analysis of a novel cardiac stimulant, UK-61 260, in human plasma by GC–MS with selected-ion monitoring 439
Uracil arabinoside
Determination of ^{3}H-labelled cytosine arabinoside and eight metabolites in cell ex-

tracts by HPLC and scintillation spectrometry 331
Uracil arabinoside phosphates
Determination of ^{3}H-labelled cytosine arabinoside and eight metabolites in cell extracts by HPLC and scintillation spectrometry 331
Vitamin B_6 vitamers
Analysis of vitamin B_6 vitamers in human tissue by cation-exchange HPLC 226
Xylazine
Rapid extraction and determination of xylazine in greyhound urine using HPLC 519